

**APPENDIX to**  
***A Systematic Review of the Job Stress  
Intervention Evaluation Literature:  
1990—2005***

LaMontagne AD, Keegel T, Louie AM, Ostry A, and Landsbergis PA (2007):  
A systematic review of the job stress intervention evaluation literature: 1990—2005.  
*Intl J Occup & Environ Health 13(3).*

**TABLE OF CONTENTS**

Appendix: Explanatory Note .....	2
Appendix Table I: Job Stress Intervention Studies with “ <i>High</i> ” Systems Approach Ratings: 1990-present.....	3
Appendix Table II: Job Stress Intervention Studies with “ <i>Moderate</i> ” or “ <i>Low</i> ” Systems Approach Ratings: 1990-present.....	21
Appendix Table III: Studies Reporting on Job Stress Intervention Evaluation Across Multiple Independent Organisations or Worksites .....	48

## EXPLANATORY NOTE: JOB STRESS INTERVENTION STUDIES 1990—2005

These tables summarise job stress intervention studies that met our specified inclusion criteria. They are summarised into three tables, each ordered alphabetically by first author (first column). The first column also includes brief description of the study population and/or setting, and the number of subjects included in the evaluation. Appendix Table I includes all studies rated as having a HIGH systems approach to job stress intervention (second column), as defined in the Methods section. Appendix Table II presents all studies rated as having a MODERATE systems approach (ordered alphabetically by first author, starting on page 21), followed by those studies rated as having a LOW systems approach (ordered alphabetically starting again within same table, by first author on page 29). Finally, Appendix Table III summarises studies of multiple worksites where varying systems level interventions were implemented (ordered alphabetically by first author, starting on page 48)

Additional notes in Systems Approach column are: intervention included employee participation (PAR); needs assessment or risk assessment conducted to tailor intervention to context (NA/RA); job stress/occupational health & safety intervention integrated with health promotion (OHS/HP). Levels of intervention (third column) are noted as physical work environment (E), organisational (O), at the interface of organisation and individual (O/I), or individual (I). Intervention duration is also noted in the third column, with indicated units ranging from hours to years. Level of causal inference (level of confidence in attributing observed effects to intervention and not other causes) was rated as follows: \*\*\* = evidence obtained without a control group or randomization but with evaluation; \*\*\*\* = evidence obtained from a properly conducted study with pre and post measures and a control group but without randomization; \*\*\*\*\* = evidence obtained from a properly conducted study with pre and post measures and a randomized control group. Additional abbreviations used include: IG (intervention group); CG (control or comparison group); WC (Workers' Compensation); Indiv (Individual); Org (Organisational); GHQ (General Health Questionnaire); SF (Short-Form). Principal findings are also summarised as favourable at the organisational level (O<sup>+</sup>), such as decrease in sickness absence, unfavourable at the organisational level (O<sup>-</sup>), or no significant difference in organisational level outcome(s) assessed (O<sup>NS</sup>), and similarly for individual-level and environmental-level outcomes. These formed the basis of summaries presented in the Figures 3 and 4 of the main paper.

## APPENDIX Table I: Job Stress Intervention Studies with “High” Systems Approach Ratings: 1990-present

Study: First Author, Year  Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High)	Intervention Level or Levels (E, O, I);  Intervention Duration	Control or Comparison Groups;  Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup>  Note if includes qualitative (QUAL)	Principal Findings
Adkins, 2000 <sup>1</sup>  US Air Force;  n =16,193	HIGH  PAR  NA/RA  OHS/HP	E, O, O/I, I  3 years (approx.)	No control group;  Org-level: suicide rates, Workers' Compensation (WC) rates, WC costs, healthcare utilization rates and healthcare costs.  Indiv-level: Measures of stress produced by personal health, threat of job loss, problems with supervisor, work relationships, conflicting responsibilities, deployment, workload, and being away from family; measures of absenteeism, work performance, and accidents; and measures of coping strategies.	3-***  QUAL	O+  Org-level: After the first year, workers' compensation rates declined by 3.9% and health care utilization rates declined by 12%. Deaths resulting from behavioural problems, including suicides, declined by 41%. From 1995 to 1996 the suicide rate decreased 38% and an additional 25% from 1996 to 1997. However, no analyses were reported on the statistical significance of these improvements.  Indiv-level (qualitative and quantitative data): Only baseline data reported ostensibly used to tailor intervention to specific contexts.
Aust, 1997 <sup>2</sup>  German bus drivers;  n= 54	HIGH  PAR	O, O/I  12 weeks	Non-intervention control group;  Measure of need for control, evaluation of the program (participation, satisfaction, perceived benefits), and positive and negative mood.	4-****	I+  Mean level of "need for control" (previously shown to predict heart disease) was significantly reduced in IG vs. CG at 12 weeks, and this effect persisted after 3 months.

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					No significant impact on mood or symptoms. Suggestions for structural changes were discussed with superiors and, after end of the group-based stress management program, they were dealt with by the official occupational health and safety committee of the company. This report restricted to assessment of immediate effects of the 12 week group program.
Barrios et al, 1997 <sup>3</sup> Managers, engineers and factory workers; n= 48	HIGH	O/I 6 months	No control group. Personal Opinion Survey (for happiness, contentment, burnout, nervousness, tension, anxiety and anger), measures of job stress, heart rate variability, and blood pressure.	3-***	O+, I+ Evaluation of an “inner quality management program” showed increases in contentment, job satisfaction, and communication, and decreases in physical symptoms and blood pressure in hypertensive individuals.
Bunce and West, 1996 <sup>4</sup> Health care workers n=202	HIGH PAR	O/I, I 3 months, 1 year	No-treatment control group; Measures of job satisfaction, motivation, health (GHQ), tension and innovation.	4-****	O+, I+ Differential impact of interventions: improvements in GHQ and satisfaction scores, and increases in innovation were experienced by PAR group.

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Cahill, 1992 <sup>5</sup> Social service employees in USA n=454 (baseline) n=313 (final)	HIGH PAR	E, O, O/I 6 months	No control group; Measures of skill discretion / development, decision latitude / authority, job satisfaction, autonomy and strain.	3-***	O+, I+ Environ level: Introduction of ergonomically designed office equipment Org-level: Implementation of a new micro-computer based information system led to significant improvement in decision latitude and job satisfaction, a structured career path was introduced for clerical staff. No change in job strain levels in the 6 month period Indiv-level: changes in skill development.
Cartwright, 2000 <sup>6</sup> UK government department employees; n=253 (experimental grp) n=90 (control grp)	HIGH NA/RA PAR	E/O, O/I Interventions 1 year and ongoing Evaluation follow-up at 1 year	Non-intervention division; Measures of well-being, job satisfaction and attitude, the Occupational Stress Indicator (OSI).	4-**** QUAL	O+, I+ Org-level: Stress levels emanating from the organizational structure and climate were significantly reduced post-intervention in the intervention group compared to the non-intervention group (no change). Significant post-intervention improvements in job satisfaction. Sickness absence due to mental disorders unchanged in intervention group (stayed at 10%), but greatly increased in the control group (10% to 30%).

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					Focus groups indicated improved organisational climate as a result of the intervention.  Indiv-level: Significant changes in 'locus of control scores' for the intervention group. No significant changes in physical and psychological health.
Eriksson et al, 1992 <sup>7</sup> Public administration employees (Sweden) n=129	HIGH PAR	E/O, O/I, I Duration not specified	Four intervention work units and a no-treatment control work unit; Measures of social support, blood lipid profiles, general health and well-being.	4-****	O+, I+  The education, discussion group, and action plan program was deemed to be more effective in groups that had a high degree of autonomy, high decision latitude and high initiative skills.  In the intervention groups there was a significant increase in "good" cholesterol HDL (high-density lipoprotein), decrease in "bad" cholesterol LDL (low-density lipoprotein), and a sharp decrease in triglyceride levels. In the intervention groups, perceptions of more stimulating work, increased feedback from supervisors, and reduced workload were reported.  In the control group, few psychosocial or physiological changes could be observed.

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Griffin, 2000 <sup>8</sup> Australian Hospital employees n=540 56% Response rate	HIGH PAR NA/RA	O, O/I, I Duration unclear. Pre-intervention survey and 2-year follow-up	No control group; Measures of organisational climate, employee morale and distress, turnover intention and non-certified sick leave.	3-***	O+, I+ Org Level: A series of paired sample t-tests showed significant improvements, across the two years of the survey, to employee ratings of leadership, professional interaction, goal congruence, recognition, decision making participation, workplace morale, workload and workplace stress. Indiv-Level: Significant improvements also observed in professional development levels and individual morale No results were reported for changes to non-certified sick leave
Griffiths et al, 2003 <sup>9</sup> UK Senior hospital nurses (H grade most senior, G grade middle seniority, F grade least senior) N = 80	HIGH PAR NA/RA	E, O, O/I 6 months	No comparison group; Baseline and follow-up survey: measures of general well-being ('worn-out' scores), overall job satisfaction, intention to leave, absence, musculo-skeletal pain, reported working conditions in terms of reported problems (e.g., lack of management support, lack of time for leave). Follow-up survey also included items on: awareness and involvement in intervention, perceived impact of the	3-*** QUAL	O+, I+ Overall, nurses reported being slightly less worn-out. There was a slight decrease in percent of G and H grade nurses intending to leave, but a slight increase in F grade nurses intending to leave. Nonetheless, both groups remained satisfied with their jobs and absence days per year remained low. Musculo-skeletal pain increased in both groups, but the reasons for this were not clear. Success rating explored amongst those

		Intervention			Evaluation		
Study: First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings		
Innstrand et al, 2004 <sup>10</sup> Staff working with persons with intellectual disabilities; n=112	HIGH PAR	E, O/I 10 months	Non-intervention control group; Maslach Burnout Inventory (MBI), measures of job satisfaction, and work stress	4-****	involved in each intervention whether working conditions or well-being had improved in comparison with those not aware of or involved in the intervention.  The increase in time allowed for administrative aspects of nurses' work, installation of computers, and appointment of housekeeping staff varied in success. The increase in administration time was successful if staffing was not an issue; however, where it was an issue its success was limited. The installation of computers if complete was viewed positively however if it was not fully completed or training had not been provided it was not recognised as a success. Study leave and training were evaluated as effective. The intervention designed to increase communication was viewed positively.	O+, I+ The intervention group showed a significant reduction in stress and exhaustion, and a strong significant rise in job satisfaction after intervention.	
Israel et al., 1989 <sup>11</sup>	HIGH	E, O, O/I	No control groups —one participatory	3-***		O+, I+	



Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
& 1992 <sup>12</sup> , and Heaney et al, 1993 <sup>13</sup> US manufacturing plant employees (86% hourly, 14% salaried) n=1100; n=176 (Heaney et al) <sup>13</sup>	PAR NA/RA	7 years (Israel et al) 5 years (Heaney et al)	action research intervention in two independent branches of one company: one with cooperative labour-management relations, the other with adversarial; Israel et al: On-going qualitative evaluation and periodic employee surveys measuring support and well-being. Heaney et al: Measures of participation, participative climate, labour /management relations, social support and depressive symptoms (CES-D).	QUAL (Heaney et al: questionnaire survey, semi-structured interviews and field observations)	Israel et al: Although there were increases in co-worker support and trust between hourly and salaried staff, job security decreased due to a down-sizing and company split during the intervention. Consistent with this down-sizing event, there was also an increase in negative feelings and sleeping problems, supervisor support, and some symptoms. Heaney et al: Employee participation in decision-making increased in both intervention conditions. Also found enhanced employee perceptions of the effectiveness of the process in both contexts, but employee well-being did not improve in either group. Labour management relations context influenced the impact of the stress project. The intervention enhanced employees' perceptions of the climate for participation only in the organization with more cooperative industrial relations. However, counter to hypothesized changes, increases in coworker support and decreases in depressive symptoms were associated with the intervention only in the organization with more adversarial industrial relations. This likely explained

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					by adversarial setting having no previous opportunities for exchanging support with co-workers. Interpretations complicated by organization restructuring during study.
Kalimo, 1999 <sup>14</sup> Forest Industry employees:  n= c.2000 respondents at 2 year follow up n=c 11,000 respondents at 10 year follow up 61% response rate	HIGH PAR NA/RA	O, O/I  Ongoing intervention with consultation over 15 years (1984-1999) including, support, training, and feedback on changes in work conditions and work behavior.  2, 4 and 10 year evaluation data collections	No control or comparison group;  Measured work-related and health-related factors: group support, commitment, and strain.	3-***	O+ I+  Org Level: Work changes viewed positively but time pressures had increased at 2 year follow up and at the 10 year follow up. Overall level of stress remained low with the majority of staff assessing their psychological working capacity as good. Job security was perceived as slightly better in the company than the general population.  Indiv Level: No differences in health indicators at 2 year follow up. At 10 year follow up workers self-evaluated their health as 'good'.
Kawakami 1997 <sup>15</sup> Japanese blue – collar manufacturing employees: n=187.	HIGH PAR (supervisors only) RA/NA	E, O, O/I 1 year	Intervention group = 2 worksites (n=79) vs comparison group from 3 worksites (n=108) with complete data at 2 years follow-up  Measures: depression symptom score, sickness absence, systolic and diastolic blood pressure, and work characteristics.	4-****	O+, I+  Statistically significant decreases in depression symptoms and days of sick leave in intervention vs control group (adjusted for gender age and baseline levels). No changes to blood pressure levels.  Percent reporting work overload increased in intervention group versus

		Intervention			Evaluation		
Study: First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>		Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings	
	Kvarnstrom, 1996 <sup>16</sup> Swedish electrical manufacturing employees; n=c50.	HIGH NA/RA PAR	O, O/I 6 years and ongoing at time of publication; 1-year follow-up compared to data 2 years prior to intervention	No control groups: Measures of production, turnover, sickness absence and workplace injuries.	3-*** QUAL		control, due to an unanticipated intervention-independent workload increase at intervention sites that did not occur at comparison sites in same company. O+, I+ Org Level: Organization re-structured from a 'taylorist' approach to the formation of 1,000 object oriented work groups. Significant reductions in turnover, absenteeism and stress-related injury rate, and a major improvement in production. Indiv Level: Individual skill levels were increased, anecdotal reports of improved quality of life.
Lavoie-Tremblay et al, 2005 <sup>17</sup> Health care workers in a long-term care unit; n=60	HIGH PAR	E, O/I 6 months intensive, 1 year follow-up	No control group; Job Content Questionnaire, Effort Reward Imbalance Questionnaire, Psychiatric Symptom Index, and an indicator for recorded absenteeism.	3-***		O+ There was a significant increase in reward and a significant decrease in Effort Reward Imbalance following the intervention. Absenteeism rates decreased from 8.26% to 1.86% over the study period, but in the rest of the institution remained the same. However, there was a significant decrease in social support from supervisors at post-test.	
Logan et al 2005 <sup>18</sup>	HIGH	E, O/I	Non-intervention control group;	5-****		O+, I <sup>NS</sup>	

Intervention		Evaluation			
Study: First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Unit managers of a trucking company n=64	PAR	10-hour training session	Measures of overall control, supervisory support, somatic complaints, depression (CES-D), anxiety, and job satisfaction (JDS).	QUAL	This was intended as a control- enhancing intervention for unit managers.  The intervention increased perceptions of control after 4 months, but only for those managers with supportive supervisors. In conjunction with supervisory support, the intervention produced improvements in job satisfaction, but not general well-being outcomes. Process evaluation interviews indicated that the intervention was implemented as intended.
Lourijssen, et al, 1999 <sup>9</sup> Employees of one Dutch hospital (n = 612) versus a control hospital (n = 382)	HIGH PAR NA/RA OHS/HP	E, O, O/I, I 3 years	Non-intervention control hospital; Org-level measures: absenteeism rates, intervention costs, and intervention benefits  Indiv-level measures: Interviews with supervisors and measures of work organisation, employee health, health behaviours and absenteeism.	4-****	O+, I+, E+  Org-level: Significant difference in absenteeism percentage in intervention versus control hospital after 3 years (4.0 versus 6.6). Greater decline over 4 years in intervention (8.9 to 4.0) than control (7.1 to 5.4) against steady rate averaged across all Dutch hospitals (6.5 to 6.6). Estimated benefits (1.6 million Guilders) exceeded costs (1.2 million Guilders) at the intervention hospital 2 years into the intervention.  Indiv-level: Improved employee opinions of how sick co-workers dealt with by management, quality of patient care, working conditions, and

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High)	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					psychosocial work climate also reported. Little self-reported impact on health behaviours, however, these programs not yet implemented.
Maes et al, 1998 <sup>20</sup> The Brabantia Project—Dutch manufacturing company employees; n = 264	HIGH PAR NA/RA OHS/HP	E, O, I 3 years	Non-intervention control group (n = 130) versus intervention (n = 134); Org-level measures: absenteeism rates Indiv-level measures: cardiovascular health risks, psychological job demands, job control, ergonomic risk factors.	4-****	O+, I+ Org-level: significant drop in sickness absence in intervention (15.8% to 7.7%) versus control (14.3% to 9.5%) groups, which by the company's determination yielded a positive financial return on its investment in the project. Indiv-level: Manufacturing employees in intervention group versus control had significantly greater favourable changes in cardiovascular health risks (decrease), psychological job demands (decrease), job control (increase), and ergonomic risks (decrease).
Matrajt, 1992 <sup>21</sup> Mexican manufacturing plant employees; n=130 managers and 3600 employees.	HIGH PAR NA/RA	O, O/I 17 weeks situation diagnosis and corrective phase, 12 months follow-up	No control groups; Measures of productivity, psychosomatic symptoms and internal relations.	3-***	E+, O+, I+ Indiv-level: Progressive reduction in psychosomatic illness (17% for managers and 15% for assembly-line workers). Org-level: General work environment improved, with an increase in productivity and reduced absenteeism. Cost-benefit evaluation justified investment in the study through

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					increased productivity, savings in lost work hours and the costs of training replacement managers, and medical care and sickness benefits for assembly-line workers.
Melchior 1996 <sup>22</sup> nurses; n=161	HIGH	E/O, O/I 1 year	No-treatment control group; Measures of Maslach burnout inventory, employee turnover	5-****	O+, I <sup>NS</sup> There was no observed change in burnout inventory scores for the treatment or control group, but job turnover decreased significantly versus controls as a result of innovation in care delivery with emphasis on primary nursing, feedback/support, and communication skills training in order to reduce burnout.
Michie, 2004 <sup>23</sup> Hospital cleaning (intervention) and catering (control) staff; n=221	HIGH PAR NA/RA	E, O, O/I Changes and new reporting systems introduced in one month Evaluation follow-up at 6 and 12 months	Cleaning staff intervention group (n=221) and catering staff control group (n=91); Org-level measure of monthly sickness absence rates	4-****	O+, I <sup>NS</sup> Significant, albeit small, reduction (2.3%) in the difference in sickness absence between intervention and control groups in the six months after intervention. The difference in sickness absence rates, however, was not maintained at 12 months.
Mikkelsen, 2000 <sup>24</sup> Norwegian Healthcare	HIGH PAR	O, O/I, I 12 week intervention.	randomized control group (n=71) non-randomised intervention group (n=64)	4-**** QUAL	O+ Org level: Improvements for work-related stress and psychological

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
employees; n=135		Evaluation 1 week & 1 year post-intervention	Measures of work stress, subjective health, demands/ control, skill discretion, decision authority, social support, role harmony, learning climate and leadership.		demands, however these may be due to differences between the control and intervention groups. Paired t-tests showed significant improvements in 'role harmony' and 'learning climate satisfaction' in the intervention group. Written reports from management, consultants and union representatives favourable regarding usefulness of intervention.  Individual level: No significant effects on subjective health and anxiety.
Mikkelsen et al, 1999 <sup>25</sup> Postal workers  N = 153	HIGH PAR	E, O 12 weeks intervention, 1 year evaluation follow-up	Compared two intervention groups (n = 91) to two control groups (n = 62); Cooper's Job Stress questionnaire Spielberger State-Trait Anxiety Inventory (STAI), Organizational Commitment Questionnaire (OCQ), Job satisfaction was measured by the Quinn and Shepard method, Job Content Questionnaire, Subjective health was measured by the Health Inventory; Social support, Learning climate and leadership were also measured.	5-*****	O <sup>NS</sup>  The goal of the intervention was to improve work environment and health, however this study was also affected by organizational restructuring and turbulence.  Work conditions deteriorated during the observation period in the control groups. In one of the intervention groups, this negative trend was reduced by the intervention. Lack of positive results in the other intervention group may have been due to organizational restructuring and turbulence.
Munz, 2001 <sup>26</sup> Customer	HIGH PAR	E, I 3 months	Four comparable work units in 4 different cities: two intervention	4-****	E+, O+, I+  Org level: 24% decrease in

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
service/sales representatives (USA) n=79		'comprehensive stress management program'  Evaluation 12 weeks after training	(combined self-management training and stressor reduction process) versus two non-intervention control units.  Org level measures: Productivity and absenteeism.  Individual measures: Pre and post intervention questionnaires, measures of perceived stress, depression, positive/negative affect.		absenteeism compared to 7% decrease in the control groups and 23% increase in productivity (measured by sales revenue) compared to 17% increase in the control group. Statistical significance of difference between intervention and controls not reported);  Indiv level: Significantly improved emotional well-being, perceived stress levels and depression.
Nijhuis et al, 1996 <sup>27</sup> Dutch construction employees; n=425	HIGH NA/RA OHS/HP	O, O/I, I  Apparently ~1 year: Interventions (organisational structuring and training) started in autumn 1992, effects expected during 1993,  Post intervention (2-year follow up) survey in 1994.	Two no-treatment control groups;  Measures of absenteeism, health complaints, and employee attitudes to work	4-****	E+, O+, I+  Baseline surveys were completed by all groups, however, authors do not specify how and if comparison groups were used in statistical analysis of pre and post data. Nevertheless, they report that: <ul style="list-style-type: none"> <li>• Significantly fewer employee complaints with respect to aspects of job content and labour relations, (p&lt;.05).</li> <li>• No significant reduction in employee complaints with respect to decision latitude, physical working conditions, stress-related fatigue, or health complaints;</li> <li>• Considerable reduction in absenteeism rates of managerial</li> </ul>



		Evaluation		
Intervention		Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration			
<b>Study:</b> <b>First Author,</b> <b>Year</b> <b>Population and</b> <b>Sample Size</b>				<b>staff;</b> <ul style="list-style-type: none"> <li>Sickness absence rates declined from 10% to 8% in the total population. Multiple regression analysis showed that 34% of this decline could be attributed to experienced differences in stressors (social relations, task information, and participation);</li> <li>Economic evaluation found cost effectiveness through reduced absenteeism of managers, greater capacity for spotting and solving problems, and improved efforts by workers.</li> </ul>
Orth-Gomer et al, 1994 <sup>28</sup> Government office workers (Sweden) n=129	HIGH PAR  O, I 8 months	Non-intervention control group (n =35) versus intervention (n = 94); Physiological measures: apolipoprotein B/apolipoprotein AI ratio Psychosocial measures: job strain and social support.	5-*****	O+, I+ Intervention included education program, relaxation training, and worker committees which developed and carried out action plans to reduce work-related sources of stress. Found: <ul style="list-style-type: none"> <li>Significant decrease in apolipoprotein B/apolipoprotein AI ratio occurred in the intervention group, but not in the control group.</li> <li>Stimulation from and autonomy over work significantly increased in the intervention group but remained the same in the control group</li> </ul>

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Poelmans, 1999 <sup>29</sup> Pharmaceutical company employees in Belgium n=324	HIGH, NA/RA	E, E/O, O/I, I 1 year. Only sickness absence post intervention evaluation completed	No control groups; 324 employees completed baseline questionnaires on stress, psychosomatic complaints and working conditions.	3-*** QUAL	O+ Environ Level: Development of purchase criteria for office furniture to meet higher ergonomic standards. Org Level: Significant decrease in absenteeism. Anecdotal feedback that occupational stress was de-stigmatized at all levels of the company.
Sastry, 1992 <sup>30</sup> Mining company in India: Managers (n = 204) Operators and loaders (n = 404)	HIGH PAR NA/RA OHS/HP	O/I, I 3 day training program	No control group; Indiv-level: measures of qualitative job content (e.g., participation in decision-making, role ambiguity, interpersonal relations), stress-related health complaints, and health behaviours	3-***	O+, I+ Reported percentages of respondents indicating improvements at first feedback period (6 months), but no numbers (response rates) nor statistical analyses reported. Examples of findings: <ul style="list-style-type: none"> <li>• 39% of the senior managers and 28% of the middle-level managers indicated improvements in qualitative content of the job</li> <li>• 47% of senior managers and 39% of middle-level managers indicated a reduction in their stress-related health complaints;</li> <li>• Shift schedules rearranged in response to need identified by operators and loaders;</li> </ul>

		Intervention			Evaluation		
Study: First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings		
Schaubroeck, 1993 <sup>31</sup> Nonacademic university employees in a Midwest USA university; n=27 in intervention group n=25 in control group	HIGH PAR NA	E, O/I 2 year	Randomized controlled trial with control group waitlisted. Org level measures: Measures of absenteeism, role ambiguity / conflict and supervisor satisfaction. Individual measures: physical and mental well- being.	5-****	<ul style="list-style-type: none"> <li>53 % of senior managers, 29% of middle-managers, 34% of operators, and 54% of loaders reported a "reduction in tobacco consumption".</li> </ul> <p>O+, I<sup>NS</sup> Org-level: ambiguity and supervisor dissatisfaction were reduced through role clarification. Indiv-level: No significant effects The study took place during a period of organizational financial cutbacks with rumours of personnel cutbacks and job reclassification that were viewed negatively by the employees</p>		
Van Dierendock et al, 1998 <sup>32</sup> Direct-care mental health professionals and staff N=149 at final	HIGH PAR	E/O, O/I, I 5 weekly group sessions of 1/2 day each Data collection pre-, 6 months post-, and 12 months post-intervention	External and internal control groups; Maslach burnout inventory, social support, turnover intention, absenteeism and equity	4-****	<p>O+, I+</p> <p>Burnout, absence, and deprived feelings (negative value in equity measure, indicating an inequitable relationship) significantly decreased relative to control groups. Turnover intention remained stable for intervention group, but increased for the</p>		

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					internal control group.

<sup>§</sup>Additional Notes: intervention included employee participation (PAR); needs assessment or risk assessment conducted to tailor intervention to context (NA/RA); job stress/occupational health & safety intervention integrated with health promotion (OHS/HP).

<sup>#</sup>3- \*\*\* = evidence obtained without a control group or randomization but with evaluation; 4- \*\*\*\* = evidence obtained from a properly conducted study with pre and post measures and a control group but without randomization; 5- \*\*\*\*\* = evidence obtained from a properly conducted study with pre and post measures and a randomized control group

## APPENDIX Table II: Job Stress Intervention Studies with “MODERATE” or “LOW” Systems Approach Ratings: 1990-present

		Intervention		Evaluation	
Study; First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Bagnara et al, 1999 <sup>33</sup> Trainee nurses n=128	MODERATE	O/I, I 6 months of discussion groups (12-15 students) plus periodic meetings with supervisor	Non intervention control group Measures of psychological well-being (GHQ), anxiety, self -esteem, work expectations and work involvement.	5-****  	O+, I+ Psychological well-being (GHQ) improved significantly within the intervention group, but improvement in relation to control group not reported. Significantly more trainees in the intervention group passed their exams in comparison to control group.
Blomkvist et al, 2005 <sup>34</sup> Coronary critical care unit nursing staff (Sweden); n=36	MODERATE	E Sound absorbing ceilings installed and data collected at start and end of work shifts to calculate daily exposure effects during baseline “sound reflecting period” of 20 weekdays, and intervention “sound absorbing” of 22 weekdays.	No control or comparison group;  Measure of acoustics, pressure (stress, calmness, hastiness), strain (irritation, anger, tension), distress (anxiety, sadness, depression) and a condensed visual analogue version of the Swedish demand-control-support model.	3-***  	O+, I+ Improved acoustics significantly reduced demands, strain, and pressure during the afternoons, while control/support and distress were not affected by the condition at all. Staff reported feeling more relaxed and less irritable during the intervention period. It was previously known that physical and psychosocial stressors can interact in producing enduring health effects. Notably, this study demonstrates that intervening on physical stressors can reduce psychosocial stress, thus exploiting the interaction in an intervention context.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Bond & Bunce, 2001 <sup>35</sup> U.K. administrative employees; n=97	MODERATE PAR RA/NA (Note: strong participatory methodology, but no indication of secondary/tertiary intervention)	E/O, O/I Five 2-hour steering committee meetings over a 3 month period, plus various activities integrated into day to day work; 12 month evaluation follow-up	Matched randomized control group; Measures of mental and physical ill- health, sickness absence (based on personnel records), performance, and job satisfaction. Used the Occupational Stress Indicator, and Job Content Questionnaire.	5-*****	O+, I+ Employee committees developed action plans to increase employee control in various areas. Significantly improved participant's mental health, sickness absence rates, and self-rated performance at 1-year follow-up Results also indicate that favourable effects mediated by increased employee job control through work re- organisation.
de Croon et al, 2004 <sup>36</sup> Dutch lorry drivers; n=78	MODERATE	E 2 years	2 matched control groups; Measures of job demands and control, mental health (need for recovery after work) and job attitudes (organisational commitment) using the Dutch Questionnaire on the Experience and Assessment of Work (VBBA).	5-*****	O- Results showed that the application of on board computer (OBC)-systems negatively affected the drivers' job control and organisational commitment. However, OBC-systems did not influence the drivers' psychological job demands and need for recovery after work. Accordingly, it was concluded that the application of OBC-systems negatively affects the lorry driver's psychosocial work environment and job attitudes.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Elo, 1998 <sup>37</sup> Finnish carton production employees; n=118 baseline	MODERATE NA/RA PAR	E/O Evaluation in one of three departments 3-years post initial survey	No control or comparison groups; Measures of variability of work, and mental and physical strenuousness (using the Occupational Stress Questionnaire).	3-*** QUAL	O+, I+ Indiv-level: Significant overall reduction in mental and physical strenuousness levels. Org-level: Significant increases in the variability of work in the department which was evaluated
Landsbergis & Vivona-Vaughan, 1995 <sup>38</sup> State government agency employees (US); n=77	MODERATE PAR	O, O/I 1 year as a pilot program	2 intervention departments and 2 waitlist control departments; Measures of communication, support, supervisor relations, job characteristics, organisational climate, job satisfaction and psychological/physical strain (i.e., depression and sleeping problems).	4-**** QUAL	O <sup>NS</sup> Mixed impact on scores of the intervention in department 2 (relative to control 2), though univariate and multivariate analyses (adjusting for demographics and job characteristics) were borderline or non-significant. Negligible or negative impact in intervention department 1 (relative to control 1), though effects were non- significant. Intervention efforts were disrupted by a major organizational restructuring and hampered by lack of formal management and labour commitment to maintaining the intervention process.
Molleman, 1995 <sup>39</sup> Healthcare employees (nurses);	MODERATE	O 8 hours per week support by a staff nurse for 6 months	Matched control groups; Measures of perceived control, autonomy and performance.	4-****	O+ The new work design brought about a shift in control from head nurses to regular staff nurses, with the following

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
n=435		6, 12 and 18 months after start of intervention			higher in intervention versus control groups: <ul style="list-style-type: none"> <li>• Level of control of nurses over primary care;</li> <li>• The extent to which nurses can make decisions autonomously</li> <li>• The necessity to consult nurses before making decisions.</li> </ul>
Parkes, 1995 <sup>40</sup> Driving Test examiners; n=49.	MODERATE	O	Cross-over control; Measuring cognitive performance under variably demanding work.	5-*****	O+ Significant decrease in both speed and accuracy while performing tasks with increasing workload.
Proctor et al, 1998 <sup>41</sup> Nursing home care staff; N=84	MODERATE	O/I, I  6 month intervention	No-treatment control group; Used the occupational stress indicator and GHQ.	4 - ****	O <sup>NS</sup> , I <sup>NS</sup> No significant differences were found between the intervention versus control group for this intervention in skill development in residential care provision.
Randall et al 2005 <sup>42</sup> Railway workers n=37 (n=25 were aware of the	MODERATE RA/NA	O Change to role responsibilities as a result of baseline needs assessment	Control group assembled post- intervention Questionnaire measurement of work stress using the General well-being Questionnaire and measurement of	3-*** (control group was assembled after the intervention from workers with	I+ Indiv level: Improvements in well-being in the group aware of the intervention compared to the groups who were not aware of the intervention.



Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
intervention, n=12 were unaware of the intervention) NOTE: Other study described in this paper reported under Griffiths 2003.		Evaluation 18 months after the risk assessment. Length of time after workers were made aware of the intervention was not specified.	exhaustion using a 12 item scale.	baseline and follow- up scores who were unaware of the intervention).	
Reynolds et al, 1997 <sup>43</sup> City council employees; n=156	MODERATE NA/RA	O/I, I 1-year and 2-year evaluation follow-up	Non-intervention control group versus individual counseling group and organizational change group; Measures of job characteristics, psychological well-being, physical symptoms, work / life satisfaction and absenteeism.	4-****	O+, I+ No changes (psychological /physical well-being or absenteeism) due to organisational change intervention aimed at increasing participation and control of employees in day to day decisions. By contrast, favourable changes at individual level were observed for individual-focused comparison group (see Reynolds [1997], LOW, in Table II below).
Rydstedt et al, 1998 <sup>44</sup> and Evans, 1999 <sup>45</sup> Swedish bus drivers n=21 questionnaire n=41 field study (observations of	MODERATE	E, O Intervention on- going over 2 years; Field study evaluation at one year; questionnaire study evaluation at 2.5 years.	Field study: 10 intervention drivers and 31 matched controls; Questionnaire for occupational stress and perceived workload; Field study for observer-rated job hassles, systolic blood pressure, heart rate at work, and perceived distress after work.	3-***	O+, I+ Intervention on a difficult inner city bus line in Stockholm--designed to reduce traffic congestion, reduce passenger service, and reduce workload demands imposed on the drivers. Investigators hypothesised that initially elevated indices of job stress among

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
workload, psychophysiological reactions at work, and self-reported stress)					<p>drivers on the difficult intervention bus route would be reduced to levels equivalent to those in comparison group: results consistent with hypothesis in field study (n = 41):</p> <ul style="list-style-type: none"> <li>• Significant decline in systolic BP (-10.7 mm Hg) in the intervention group greater than the comparison group (-4.3 mm Hg);</li> <li>• Significant decline in heart rate (3.7 bpm) in the intervention group greater than the comparison group (0.5 bpm);</li> <li>• Significant decline in job hassles per hour (-4.5) in the intervention group greater than the comparison group (+0.6);</li> <li>• Changes in job hassles were significantly correlated to changes in systolic blood pressure, heart rate, and perceived stress, but not diastolic blood pressure change</li> </ul> <p>This further supported by favourable decreases in perceived workload and distress in the smaller questionnaire study (n = 21)</p>
Smith et al, 1992 <sup>46</sup>  US meat	MODERATE PAR NA/RA	E, O 1 year	No control group; Unstructured interviews and conversations including discussion of	3-***  QUAL	O <sup>NS</sup>  Overall, there seems to be some improvement in psychological

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
processing employees;  N =~ 125-150 (not specified)			health status, psychosocial (e.g., job stress, job satisfaction), and other working conditions		indicators (nervous or irritable) for meat cutters and for meat wrappers, but not for meat processors. For psychosocial factors there appears to be an overall worsening effect for meat cutters and meat processors.  Interviews showed that meat processors and meat wrappers had very positive feelings about the job enlargement program. They reported that their overall job satisfaction was greatly increased, even though this was not reflected in their responses in the survey. Meat cutters were generally happy with increased rotation away from cutting meat, but unhappy with the lower job content level of meat processing and wrapping. All groups felt that there was less job stress overall.
Smoot & Gonzales, 1995 <sup>47</sup>  State hospital employees; Intervention n=35 Control n=37	MODERATE	O/I, I  4 weekly 8-hour sessions, small group (6-8) intensive training on work time including role play, skill development and feedback to change strategy of managing mental	Waitlist control group;  Equivalence of baseline staff characteristics between two units was confirmed (years of experience, daily hours patient contact, emotional exhaustion using the Maslach Burnout Inventory and relationships dimension using the Ward Atmospheric Scale).  Measures of turnover, sick leave,	4-****	O+  Org-level: Although no statistical testing performed, this “empathic skills training” intervention evaluation showed (in terms of % change over time): <ul style="list-style-type: none"> <li>• Reduced staff turnover in IG (increased in CG)</li> <li>• Larger reduction in sick leave in the IG than in the CG</li> </ul>

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
		health patients. Evaluation 6 months pre- and 6 months post-intervention. Essentially a job re- design implementation.	annual leave, patients' rights complaints, assaults of staff, and cost- benefit analysis.		<ul style="list-style-type: none"> <li>Reduced annual leave in IG (increased in CG)</li> <li>Reduction in patients' rights complaints filed (with increase in CG)</li> <li>Smaller reduction in assaults on staff in the IG than the CG</li> </ul> <p>Cost-benefit analysis revealed substantial savings for the trained unit and increased expenditures for the control unit, despite the high cost of time required for this intensive training.</p>
Terra, 1995 <sup>48</sup> Dutch metal can manufacturing plant employees; n=430.	MODERATE PAR NA/RA (physical work environment) Note: Participation level very high, with workers involved in job redesign.	E, O, O/I Initial intervention period not clearly specified- apparently 6 months, 5 years of follow up	No control group; Org level measures: Measures of productivity, sickness absence rates.	3-***	O+ Org-level: <ul style="list-style-type: none"> <li>50% reduction in sickness absence rate in comparison to significant increases in comparable plants (as reported by company OHS Service), saving 1 million Guilders per year;</li> <li>66% increase in productivity, from 0.26 to 0.43 million cans/worker/year.</li> </ul> Indiv-level: not systematically assessed. Managers anecdotally reported that workers better qualified, informed, and motivated.
Theorell 2001 <sup>49</sup>	MODERATE	O, O/I	No treatment control group versus	4-****	O+, I+

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Swedish Insurance company employees; n=299		2hr biweekly training for managers over 1 yr  One year follow up	intervention with baseline and one year follow-up; Baseline and one year follow up. Blood sampling for cortisol levels (baseline n=153 intervention, n=146 control). Questionnaire demand control model and work climate (baseline n=119 intervention, n=132 control).		Org Level: Statistically significantly greater improvement in decision authority post intervention for employees via management training.  Indiv Level: Statistically significant decrease in serum cortisol levels (employee arousal) post-intervention
Wahlstedt & Edling, 1997 <sup>50</sup> Swedish postal employees; n=100.	MODERATE	E/O, O/I, O Changes to ongoing systems and structures introduced in one month	No comparison groups; Measures of psychosocial factors' sleep disturbances, gastrointestinal complaints and sick leave.	3-***	O+, I+ At one-year follow up, significant increase in skill discretion and perceived authority was significantly correlated with lower levels for sleep difficulties and gastrointestinal complaints.
Alexander, 1993 <sup>51</sup> White collar employees in automotive industry; n=86	LOW OHS/HP	I	Matched controls; Measured skin conductance, general health, trait anxiety, work tension, sleep problems and job satisfaction.	4-****	I+ Significant improvements in reducing skin conductance, trait anxiety and alcohol/cigarette use in comparison to the control group for regular attendees of MED program. Less effect for

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					irregular attendees.
Bond et al, 2000 <sup>52</sup> Media organization employees; n=90	LOW	I 9 hrs (over 3 months)	Randomized control group (waitlist, n = 30), 'emotion-focused' coping skills group (n = 30), and 'problem-focused' innovation promotion program group (n = 30);  Measured GHQ, depression, motivation, job satisfaction and attitudes toward innovation and change.	5-*****	I+  Indiv level: Improvements in mental health and work-related variables were found in both intervention groups (two types of stress management intervention). As hypothesized, changes in outcome variables in the coping skills group were mediated only by the acceptance of undesirable thoughts and feelings. In the 'problem-focused' Innovation Promotion Program, change was mediated only by attempts to modify stressors.
Carson et al, 1999 <sup>53</sup> Mental health nurses; n=53	LOW	O/I  Intervention duration unclear	Standard-care control group (provided with booklet on stress management) versus intervention group (individual feedback and support). Both groups received some form of feedback on questionnaire scores;  Measures included the DeVillers Carson Leary Stress scale, GHQ, Maslach burnout inventory.	5-*****	I-  Unexpectedly, greater stress reduction (in Stress Scale scores) was indicated in the control group (MD = -11.5) rather than the intervention group (MD = -3.0). These differences were statistically significant.  In addition, this finding was supported by similar patterns in the related measures of GHQ and Maslach Burnout Inventory. Attrition of study subjects may have affected the results.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Cecil, 1990 <sup>54</sup> School teachers; n=54	LOW	O/I, I	Co-worker support, stress inoculation training, or no-treatment control; Measuring teacher stress.	5-*****	I+ Stress inoculation training was effective in reducing teachers' self-reported stress, while the co-worker support group was not
Cigrang, 2000 <sup>55</sup> Military trainees; n=178	LOW	I	"Usual care" control group: Measured graduation /discharge rates.	5-*****	O <sup>NS</sup> No significant results reported after stress inoculation training.
Cooper, 1992 <sup>56</sup> English Postal employees; N=288	LOW NA/RA	I 2 years	Control group (non randomized) Org level measures: Measures of absenteeism Individual measures: mental health, self-esteem, organizational commitment and changes in health behaviors	4-****	O+, I+ Org-level: Significant improvement in absence rates. Indiv-level: Decline in anxiety levels and depression, and increase in self-esteem. However, no marked changes to satisfaction and commitment levels.
Delvaux et al, 2004 <sup>57</sup> Oncology nurses; n=115	LOW	O/I 3 weeks (105 hours)	Non-intervention control group; Measures of nurse attitudes, communication skills, and occupational stress levels.	5-*****	I+ Compared to controls, nurses who participated in a psychological training program (PTP) reported positive changes on their stress levels, communication skills, and attitudes.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Doctor, 1994 <sup>58</sup> UK Police Officers N=61	LOW	I	Non-treatment control group Intervention 12 week group counselling sessions GHQ measures, 19 item occupation- specific stress situation questionnaire and measures of absenteeism.	5-*****	O <sup>NS</sup> , I <sup>NS</sup> Indiv Level: No significant differences between control and intervention groups, nor between baseline and post intervention scores. Treatment group participants provided feedback that sources of perceived stress were related to intrinsic factors within the organization. Org Level: No differences on levels of sickness absence.
Elliott, 1991 <sup>59</sup> US pharmaceutical employees; n=56	LOW NA/RA	I Four 2-day workshops	No control group: Hassles Scale administered by mail one-month after workshops) and Myers Briggs Type Indicator.	3-*** QUAL	I+ After-only: Positive subjective evaluations from participants of the relevance/usefulness of program. Indiv-level: One month after, found lower Hassle Scale scores in intervention group (but with 42% response rate).
Eriksen et al, 2002 <sup>60</sup> Postal service employees (Norway); n=860	LOW OSH/HP	O/I 12 weeks	Non-intervention control group (n =344), physical exercise only (n = 189), and integrated physical exercise and stress management training (n – 162); Subjective Health Complaint Inventory (SHC), self-reported sick leave,	5-*****	O <sup>NS</sup> , I+ The exercise-only group showed improved general health, physical fitness and muscle pain, and the integrated exercise and stress management group showed improved stress management, with the



Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
			Cooper job stress questionnaire.		integrated group showing the strongest effects. Notably, however, there were no significant effects on subjective health complaints, sick leave, or job stress.
Ewers et al, 2002 <sup>61</sup> Forensic mental health nurses; N = 33	LOW	O/I, I	Maslach Burnout inventory	5-****	I+ Significant improvements to the nurses' knowledge and attitudes towards clients.
Francis, 1992 <sup>62</sup> University employees n=43	LOW	I	Other activity group who wrote about nontraumatic events; Measuring blood samples, absenteeism, positive/negative affect and emotional inhibition.	5-****	O+, I+ Positive trends showing improvement absenteeism and in blood values (except cholesterol) in intervention group. No substantial differences in well-being between the intervention and control groups.
Freedy, 1994 <sup>63</sup> US nurses; n=87	LOW	I 5 weekly 75 minute sessions	Lagged intervention control group;  Dual Resource Intervention (DRI) (targets both social support and mastery) treatment group and Single Resource Intervention (SRI) delayed comparison group (also served as no treatment control);	4-****	O+, I+ DRI group reported significant improvements in social support and mastery compared to the no treatment control, which persisted through a 5-week follow-up. DRI participants with low initial levels of social support or mastery reported significant reductions in psychological distress. SRI group

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
			Measures of social support, mastery of destiny, emotional exhaustion, depression (CES-D) and conservation of resources.		reported a slight improvement in mastery compared to the no treatment control group.
Gardiner et al, 2004 <sup>64</sup> General Practitioners; n=105	LOW	O/I 15 hours	Non-intervention control group; Measures of work-related distress and morale, quality of work-life, and general psychological distress (GHQ).	4-****	I+ Following this cognitive behavioural stress management training program, GPs' quality of work life and morale improved while their work-related distress and general psychological distress decreased.
Goodspeed, 1990 <sup>65</sup> Various occupations within a worksite n=148	LOW	I Five 90-minute workshops for "Time-Life" and two 4-hour sessions for Myers-Briggs. Evaluation follow-up 6 to 8 months after baseline.	Time-Life Stress Management Program group (n=113) and Myers-Briggs program (n=35); (n=91 completed baseline and follow-up questionnaires). Used Stress Potential Survey (however no results reported) and Strain Questionnaire (measures physical, behavioural, and cognitive symptoms).	3-***	I+ Baseline strain scores for the Time-Life group were significantly higher than for Myers-Briggs group, further complicating interpretability of evaluation (in addition to imbalance in group sizes, and having no control). Significant reductions in follow-up symptoms measured in each of the three sub-scores as a result of both programs, although no between-group differences were identified at follow-up.
Greco, 1992 <sup>66</sup>	LOW	O/I, I	No control group	3-***	I+

		Intervention		Evaluation	
Study; First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Canadian Government employees, n=377 managers and n=1,020 employees completed training	NA/RA		Duration of intervention unspecified. Measurement of health, interpersonal relations and general climate at work. n=229 completed baseline questionnaire Unspecified number completed the evaluation		Individual level: Although no formal statistical evaluation between baseline and follow up measures was presented, managers reported that new active files on employees with functional problems had fallen by half. Managers also reported an improved capacity for action. Also, both managers and employees reported improvements to their ability to control their stress levels. Employees also reported using new techniques to manage stress and the ability to prevent stressful situations at work.
Grossman, 1993 <sup>67</sup> Healthcare professionals n=41	LOW	O/I	No control groups: Measuring the effectiveness of support group.	3-***	I+ Support groups experienced high drop out rates (perhaps individuals who needed the most help), however, participants of the program reported stress alleviation.
Heron, 1999 <sup>68</sup> UK Pharmaceutical employees; n=608. (n=368	LOW	O/I, I Stress management workshop (duration unspecified)	Control group; Pre & post intervention: GHQ, Post- intervention only: Coping skills, stress management awareness and life events questionnaire. Evaluation data	4-****	I+ Indiv Level: non-intervention group less likely to have an understanding of the principles of the management of stress in staff and more likely to have poor

		Intervention		Evaluation	
Study; First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
intervention n=240 control)			collection at varying times post intervention		coping skills. Short term improvements in self-rated well-being were not sustained.
Hyman , 1993 <sup>69</sup> Long-term care facility employees; N=51	LOW	O/I, I Three 3- hour sessions designed to address team building, communication skills, self-esteem, and stress management; Evaluation at session end	No control groups; Human Services Survey (Maslach and Jackson, 1981) (including emotional exhaustion, depersonalizations, and personal accomplishment and measures of work atmosphere.	3-*** QUAL	I+ Using a retrospective pretest design (i.e., after-only), a statistically significant (n=42) improvement from "then" to "today" was found for: <ul style="list-style-type: none"> <li>• Depersonalization</li> <li>• Emotional Exhaustion, and</li> <li>• Personal Accomplishment.</li> </ul> Responses to an open-ended question about workshop effects corroborated the quantitative data. Open-ended interview question: participants reported an increase in self-esteem, improved communication, enhanced coping skills to deal with stress and an improvement in work atmosphere.
Iwi et al, 1998 <sup>70</sup> Local authority Housing Department employees	LOW	I 3 months	Non-randomized control groups; General Health Questionnaire (GHQ) and Occupational Stress Indicator (OSI)	4-**** INS	Workers accepting the offer of counseling were subject to greater levels of work stress, had poorer self- reported health and markedly lower

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
n=193					levels of job satisfaction than those who did not. Questionnaire scores were not significantly different before and after counseling, giving no evidence of treatment effects on symptomatology. However, almost all subjects rated counseling as having been extremely helpful. This study suggests that adverse effects on staff facing organizational change may be ameliorated by improved management practice.
Jenkins, 1991 <sup>71</sup> Female public school teachers, Georgia, USA n=124	LOW	I Three hour seminar with or without individualised stress management plan	Compared seminar with step-by-step individualized stress management plan (intervention) versus global comparison (seminar without individualized stress management plan);  Questionnaire items on types of stress experienced, sources of stress at work and home, effects of stress and burnout, variety and type of stress relief methods used, and degree of involvement.	3-***	I+  Three weeks after the training, teachers provided with individualized step-by-step training reported a significantly greater increase in time spent managing stress versus the comparison group.
Johanning, 1996 <sup>72</sup> Munich mass transit operators;	LOW OSH/HP	I One year, bi-weekly 7½-hour program sessions	Waitlist control group;  Measures of musculoskeletal problems, CVD risk factors (Rose questionnaire), electrocardiograms,	4-****  QUAL	I+  Indiv-level: Intervention group overall CVD risk reduced but not significantly compared to control group.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Intervention n=98 Controls n=26		Evaluation after 1 year	HDL cholesterol, total cholesterol, and systolic blood pressure. Psychological profile ("sense-of-life") based on in- depth interview.		Intervention group back problems were significantly reduced compared to controls. Org-level: No significant difference with respect to job strain.
Kushnir, 1993 <sup>73</sup> Safety officers; n=40.	LOW	I Five weekly meetings	No-treatment control group: Measures of cognitive weariness, somatic complaints, irrational beliefs and assertiveness.	4-****	I+ Assertiveness, somatic complaints, and irrationality improved in the short term, and to a lesser extent 18 months later. Cognitive stress symptoms decreased in the long term. Assertiveness was improved in the short term.
Le Scanff et al, 2002 <sup>74</sup> Male police officers n=150	LOW	O/I 6 months: initial four-day meeting, a two-day follow-up meeting one month later, and a final one-day meeting at six months).	No control group; A stress manifestations inventory (Adaptability Questionnaire, ADQ), group interviews/discussions.	3-*** QUAL	This essentially a qualitative process evaluation. Results indicated that the psychological training was very well received and led the police management to consider contributing factors and manifestations of stress in a more extensive way. However, not able to assess impacts on stress manifestations, as most participants kept for themselves their ADQ scores. Subjective comments and feedback from the participants formed the basis of the program evaluation.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Lee et al, 1994 <sup>75</sup> Nurses; n=57	LOW	I  Intervention duration unclear	Placebo intervention.  Used the perceived stress scale.	5-*****	I+  Findings indicated a greater decrease in stress for the treatment group as compared to the placebo group.
McCue et al, 1991 <sup>76</sup> Physicians; n=64	LOW	O/I, I  ½ day workshop  Evaluation 2 weeks pre-intervention and 6 weeks post.	Non-intervention comparison group (interested volunteers who could not be freed from clinical duties);  Measuring burnout, stressors, stress symptoms and support skills.	4-****	I+  This “modest, inexpensive stress management workshop” showed positive impacts of learning and practicing interpersonal skills that may increase the availability of social support;  Intervention group reported a reduction in burnout levels and stress symptoms, and reported being more aware of work stressors and of support seeking opportunities.
Meier 2000 <sup>77</sup> Social workers; n=52	LOW	O/I  10 weeks	Non-intervention control group;  Occupational Stress Inventory	5-*****  QUAL	I <sup>NS</sup>  No statistically significant changes in levels of occupational stress or psychological strain. The small sample size and relatively weak intervention of this feasibility study made it unlikely that any effects would be detected.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Michie, 1996 <sup>78</sup> UK Hospital staff; n=92 used the service more than once n=41 participated in 6 month evaluation	LOW	I Intervention at the most 3 x 1 hr sessions Evaluation 6 months post intervention	No control group; Eight item questionnaire regarding self-assessed levels of anxiety, depression, work functioning, and work satisfaction. Workers were asked “do you feel anxious, do you feel depressed” etc. Questionnaires were completed at the beginning of the first session, at the end of the last session and six months after the last session.	3-***	O+, I+ Indiv level: For the 41 staff who completed the six month questionnaire there were significant improvements for responses to questions regarding anxiety, depression, work satisfaction, life satisfaction and functioning at work. Org Level: Sickness absence was significantly reduced amongst the 92 people using the service more than once.
Pelletier et al, 1999 <sup>79</sup> Bank employees; n=136	LOW	I 6 months	Mail, mail plus telephone, and control group; Measures of job strain (JCQ), objective wellness (Stanford SMART health-risk appraisal), perceived wellness (Brief Symptom Inventory), self-efficacy, and feelings about personal control.	5-*****	O <sup>NS</sup> , I+ No significant differences among groups were found in changes of any scales in the JCQ. At 1-year follow-up, the telephone group showed the largest improvement in mental health status rating, followed by the mail group, with a significant difference between the phone group and the control group. The telephone group showed increases in self-efficacy, perceived wellness, and feelings of personal control, with significant differences between the phone group and the control group. At 6-month assessment, the telephone group showed significant decreases in somatization and anxiety, but these



Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					differences were no longer evident at 1-year follow-up.
Peters, 1999 <sup>80</sup> Maintenance workers in Hawaii, USA n=50.	LOW OHS/HP	I Ten weeks	Control group (wait listed);  Indiv measures: Physical (e.g., blood pressure, cholesterol, overweight) and behavioural measures (exercise, smoking in health risk appraisal), self efficacy, health locus of control, state-trait personality, health attitudes and behaviour.  Org level: rates of injury, absenteeism, overall measure of job morale, satisfaction, and productivity.	5-*****	O <sup>NS</sup> , I+  Indiv level: Statistically significant Improvements in IG versus CG on: <ul style="list-style-type: none"> <li>• health risk appraisal (more health behaviour changes),</li> <li>• number of people who lost weight in IG versus CG;</li> <li>• in self-efficacy for stress management, exercise, and nutrition.</li> </ul> There were no significant effects on emotional or psychological variables with the exception of "curiosity"  Org level: No significant effects on rates of injury or absenteeism, or overall measure of job morale, satisfaction, and productivity.
Pruitt, 1992 <sup>81</sup> Government employees; n=64.	LOW	I.	Waitlist control group;  Measuring blood pressure, psychiatric symptoms, anxiety and life events.	5-*****	I+  Significant reduction in reported stress-related physical symptoms. No major effect on anxiety and blood pressure.
Reynolds, 1997 <sup>43</sup>	LOW	O/I, I	Non-intervention control group,	4-****	O <sup>NS</sup> , I+

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
City council employees; n=156.	NA/RA	Three 1-hour counselling appointments; 1-year and 2-year evaluation follow-up	individual counseling group, and organizational change group; Measures of job characteristics, psychological well-being, physical symptoms, work / life satisfaction and absenteeism.		Individual counseling intervention improved the physical and psychological well-being of employees. There were no significant differences in absence from work after the intervention was introduced.  Repeated measures MANOVA indicated that there were nonsignificant effects of comparison group and of time, but that there was a significant time X comparison group interaction [F(4,160)=2.45,p<.049], suggesting there were differential changes in the three comparison groups (see Reynolds [1997] in moderate table).
Reynolds, 1993 <sup>82</sup> Female health service employees; n=92	LOW	I Six 2-hour stress management workshops at weekly intervals	No control group; Work / life satisfaction, general health questionnaire (GHQ), session evaluation and session impact.	3-***	O <sup>NS</sup> , I+ Significant reductions in psychological distress (GHQ scores), but no changes in job or non-job satisfaction.
Robinson, 1993 <sup>83</sup> Emergency service, welfare and hospital employees;	LOW	I 11 debriefings for welfare agency employees, 18 for emergency service	No control group; Measuring impact of actual incident, stress symptoms and value of debriefings.	3-*** QUAL	I+ Employees who reported symptoms of stress following critical incident also reported these to be reduced as a consequence of psychological

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
n=172		personnel, and 2 for nurses  2 weeks post- intervention evaluation			debriefing (96% of emergency service workers, and 77% of welfare/hospital staff).  Personal descriptions of how the debriefing was helpful were reported by 75% of emergency service personnel and 84%. The debriefing was valued more by staff who were more severely impacted.
Schaufeli, 1995 <sup>84</sup> Dutch Community nurses; n=64.	LOW	I  3 day workshop  Evaluation by self- administered questionnaire one month after the workshop.	No control groups;  The Maslach Burnout Inventory and the 21 item 'Tedium' measure were used to assess the nurses' level of burnout. Reactivity was measured by the 22 item Strelau's Temperament Inventory.	3-***	I+  Workshop participants who were classified as 'low reactive' showed significant reductions in emotional exhaustion and psychological strain. No significant changes were observed for workshop participants classified as 'high reactive'.
Sheppard, 1997 <sup>85</sup> High security US government agency employees; n=44.	LOW	I  For both groups, 5 hours of instruction and 12 one-hour bi- weekly group meetings. Evaluation at the	Transcendental meditation (TM) treatment group (n=22) and corporate stress management (CSM) education group (n=22).  Blood pressure and questionnaire analyses (State anxiety, trait anxiety, depression and self-concept).	5-*****	I+  Indiv level: Trait anxiety and depression levels were significantly reduced in TM compared to CSM group. Three years post intervention, significant improvements were observed for TM compared to the CSM group in state/trait anxiety, depression and self-perception.

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
		end of the 12 weeks and 3 years post intervention			No differences found in blood pressure for either group.
Taylor, 1991 <sup>86</sup> Nurses; n=102	LOW	I	Had treatment control group with random allocation. Measures of perceived stress scale	4-****	I+ Significant difference between the control and treatment groups in stress reduction.
Teasdale, 2000 <sup>87</sup> Pharmaceutical company employees; n=452	LOW OHS/HP	O/I, I Workshops (duration not reported) ran over a 6-year period (subjects had attended at least one)	Non-intervention control groups Measures of well-being (GHQ), coping skills, life-events and stress awareness.	3-*** (after-only)	I <sup>NS</sup> Indiv- level: Although GHQ scores fell 10% 2-3 months following the workshop, these returned to their original levels 1-8 years after the intervention. Workshop attendees only had slightly better coping strategy scores than non-attendees.
Toivanen, 1993a <sup>88</sup> ; Toivanen, 1993b <sup>89</sup> Finnish Hospital cleaners and bank employees N=98	LOW	I 6 months intervention Evaluation at baseline, 3 & 6 months concurrent with intervention.	Randomized control group; Individual measures of cardiovascular ANS function. Measures of sickness absence were collected for the hospital cleaners. At the end of the intervention, interviews regarding the workplace were conducted with the bank employees, and hospital employees	5-***** QUAL	I+ Indiv level: Regular daily practice of relaxation methods normalized cardiac ANS function. Better results were noted amongst participants where relaxation practice took place as a workplace group activity than for individuals practicing on their own. Org level: Amongst the hospital

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
			completed a questionnaire on occupational factors and perceived personal health.		cleaners sickness absence decreased in both the control and the intervention groups.
Tsai, 1993 <sup>90</sup> Nurses (Taiwan); n=137.	LOW	I  Intervention included 3 relaxation training sessions at weeks 1, 2, and 5  Evaluation at weeks 2 and 5.	Non-intervention control group;  Measures of Nurses' Stress checklist (NSC) and Chinese General Health Questionnaire (CGHQ).	5-*****	I+  Intervention group reported a reduction in stress levels and symptoms after completing training course: <ul style="list-style-type: none"> <li>• Mean scores for both NSC and CGHQ differed significantly between intervention and control groups at 5 week follow-up.</li> <li>• CGHQ scores also differed significantly at 2 week follow-up.</li> </ul>
Vines, 1994 <sup>91</sup> Unspecified workers from 4 corporations (USA); n=68	LOW	I  6 training sessions over 9 weeks intervention.  Baseline 3 weeks prior to intervention, Evaluation at 20 weeks.	Non-randomized waitlist control group. (n=37) waitlist control vs (n=31) intervention.  Measures of psychological distress and health seeking behaviors.	4-****  QUAL	I <sup>NS</sup>  No improvement in the intervention group for psychological distress or health seeking behaviors.
Whatmore, 1999 <sup>92</sup>	LOW	I  Intervention period: 1 hour information	Waitlist (n=52) and non-volunteer (n=79) control groups versus 'personal stress awareness,' (n=48) 'exercise,'	5-*****	O <sup>NS</sup> , I+  Indiv level: the exercise intervention group reported significant

Study; First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Public sector employees (UK); n=270.		sessions plus 2-hour workshops on 3 different topics followed by take- home assignments.  Evaluation follow-up at 3 and 6 months	(n=43) and 'cognitive restructuring' (n=48) intervention groups;  Measures of anxiety, depression, mental and physical well-being, organisational commitment, job satisfaction and self-reported sickness absence. Questionnaire assessment of organizational commitment, job satisfaction and self-reported sickness absence		improvements for anxiety, depression, mental and physical health, and the awareness group showed improvements in depression, physical and mental health at the 3 month post- intervention stage. There were no improvements for the cognitive group. After 6 months the only significant result was a difference in somatic anxiety in the exercise group, however there was also a significant decrease in the control group.  No "significant effects" (data not shown) on job satisfaction or organisational commitment.  Org Level: Sickness absence increased in all groups with the exception of the 'exercise' intervention group, which showed a decreased sickness absence rates 6-months post- intervention (no statistical analysis or comparison of difference with controls provided).
Wiholm 2000 <sup>93</sup> Swedish Software developers working with computers n=106	LOW	I  1-1.5 hours of training over a three month period	Control group, one department selected for the intervention compared to another department without the intervention.  Individual measures: Baseline and post-training questionnaires. Blood	4-****	I+  Indiv-level: Study assessed the effects of stress management training on skin symptoms. Stress management training was associated with a significant decrease in skin symptoms

Study; First Author, Year  Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High)  Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I);  Intervention Duration	Control or Comparison Groups;  Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup>  Note if includes qualitative (QUAL)	Principal Findings
			samples were taken at baseline, immediately post –training and 5 months post-training. Blood level testosterone, cortisol and prolactin were measured.		only during the actual training period. No beneficial effects measured 6 months post-training.

<sup>§</sup>Additional Notes: intervention included employee participation (PAR); needs assessment or risk assessment conducted to tailor intervention to context (NA/RA); job stress/occupational health & safety intervention integrated with health promotion (OHS/HP).

<sup>#</sup>3- \*\*\* = evidence obtained without a control group or randomization but with evaluation; 4-\*\*\*\* = evidence obtained from a properly conducted study with pre and post measures and a control group but without randomization; 5-\*\*\*\*\* = evidence obtained from a properly conducted study with pre and post measures and a randomized control group

**APPENDIX Table III: Studies Reporting on Job Stress Intervention Evaluation Across Multiple Independent Organisations or Worksites**

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
Eklof et al, 2004a&b <sup>94, 95</sup>  White-collar computer users: 40 groups from 11 private and public organizations (Sweden);  n=342	Varying levels of Systems Approach—internal comparisons  Included NA/RA and PAR to varying degrees.	E, O, O/I to varying degrees by organisation.  Evaluation: baseline and 6-month follow-up employee surveys	Internal comparisons among the work groups, with data aggregated to work group level (n = 40);  Measures of characteristics of change processes by degree of:  • Employee participation (and empowerment) in efforts to improve work environment;  • Integration of work environment issues with traditional core organisational issues;  Were related to work environment and health improvement indicators:  • Job control, psychological demands, social support, emotional stress, comfort during computer work, and physical complaints.	3-***	This was essentially a process evaluation study, assessing the degree to which the characteristics of the change process was related to intervention-associated improvements in ergonomic and psychosocial work environment.  The change process characteristics of employee participation and integration of OHS with core organisation business were highly correlated, and may together constitute a positive organisational “learning strategy for change”.  At follow-up and prospectively, high employee participation was consistently associated with:  • lower demands (-)  • higher social support (+), and  • less stress (-)  At follow-up and prospectively, high degree of integration was consistently associated with:



Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					<ul style="list-style-type: none"> <li>• lower demands</li> <li>• higher social support (+), and</li> <li>• less stress (-).</li> </ul> <p>Neither participation nor integration was consistently associated with job control, quality of work environment modifications, comfort, or prevalence of musculoskeletal complaints.</p> <p>These findings support the importance of worker participation and integration of OHS into core organisational concerns to enable and facilitate work environment improvements.</p>
Lindstrom 2000 <sup>96</sup>  Employees from 217 small and medium-sized enterprises in manufacturing, traffic, service, and office work sectors (Finland);  n=4068  217 workplaces	Varying levels of Systems Approach—internal comparisons  NA/RA: each enterprise and employee received feedback on baseline survey.	E, O, O/I, to varying degrees by organisation.  Interventions up to 2 years in duration  1-2 years intervention and evaluation timeline	Internal comparisons between workplaces (n = 217);  Surveys done on both employers and employees pre- and post-intervention.  Measures of employee well-being (psychological strain), job satisfaction, sickness absence in previous 12 months.  Organisational practices and climate assessed in terms of : <ul style="list-style-type: none"> <li>• co-worker relations</li> </ul>	3-***	For all worksites combined, over the intervention period there was a decrease in perceived physical work environment risk factors and workload, and an increase in job control. Time pressure, however, had increased over the intervention period.  Based on employers' reporting, the statistically significant effects of interventions were most pronounced on organisational climate: <ul style="list-style-type: none"> <li>• A collaborative/ participatory approach applied in the intervention</li> </ul>

Intervention		Evaluation	
Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)
<p><b>Study:</b> First Author, Year Population and Sample Size</p> <p>PAR to varying degrees by workplace.</p> <p>Pooled analysis of a centrally coordinated intervention project, wherein participating worksites tailored intervention to their needs.</p>		<p>supervisory support</p> <ul style="list-style-type: none"> <li>continuous improvement practices</li> <li>informing about changes</li> <li>future insecurity of job</li> <li>appreciation of one's work;</li> </ul> <p>Planned organizational interventions were also characterized by focus:</p> <ul style="list-style-type: none"> <li>Customer service</li> <li>Multi-skilling</li> <li>Leadership</li> <li>Collaboration between employees and managers (~employee participation)</li> </ul> <p>Organisational effectiveness was assessed in terms of (manager perceptions on 100-point scales):</p> <ul style="list-style-type: none"> <li>Productivity</li> <li>Profitability</li> </ul>	<p><b>Principal Findings</b></p> <p>correlated with many changes in organisational climate, and most of all with an increase in continuous improvement practices (<math>r=0.36</math>, <math>p&lt;0.001</math>).</p> <ul style="list-style-type: none"> <li>Collaboration was also related to improved co-worker relations (<math>p&lt;0.05</math>) and positive changes in job future insecurity (<math>p&lt;0.05</math>), while leadership interventions decreased job insecurity less than other interventions (<math>p&lt;0.05</math>)</li> <li>Organisations with more intensive planned interventions in customer service (<math>p&lt;0.05</math>) and collaboration (<math>p&lt;0.05</math>) had a smaller increase in sickness absence compared with organisations having minor or no intervention;</li> <li>Job satisfaction did not decrease (<math>p&lt;0.05</math>), if the company carried out interventions in collaboration, "as compared to the others"</li> </ul> <p>Based on employees' reporting:</p> <ul style="list-style-type: none"> <li>All intervention types were all related an increase in continuous improvement (<math>p&lt;0.001</math>), in informing workers about changes (<math>p&lt;0.05</math>), and in appreciation of one's work</li> </ul>

Study: First Author, Year Population and Sample Size	Intervention		Evaluation		
	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
					<p>(p&lt;0.01)</p> <ul style="list-style-type: none"> <li>• Work capacity improved somewhat along with the multi-skilling of workers (r=0.06, p&lt;.01)</li> <li>• Exhaustion symptoms decreased when the interventions were focused at collaboration (r=0.06, p&lt;.0.001)</li> </ul> <p>Good profitability was related statistically significantly to all types of organisational development interventions. High production was related only to interventions dealing with multi-skilling of workers and the development of managerial practices.</p> <p>Results overall support a healthy work organisation model wherein the well-being of employees is central to company effectiveness.</p>
Nielsen et al, 2002a and 2002b <sup>97, 98</sup> Pharmaceutical, technical services, and nursing home employees at various worksites (Denmark);	Varying levels of Systems Approach—internal comparisons.  Includes NA/RA	E, O, O/I—varying degrees.  3 years  Post-intervention surveys 2 and 5 years after baseline	High-absence intervention work-sites (n = 22), high-absence control work-sites, and low-absence control work-sites (n = 30 control sites)—comparisons apparently among individual worksites (unclear).  Job Content Questionnaire, SF-36, scales on meaning of work and	4-****	At baseline, absence days per annum was significantly and positively associated all stress indicators, and was significantly and negatively associated with general health, vitality, and mental health.  Improvements were achieved, but to very different degrees in different

		Intervention		Evaluation	
Study: First Author, Year Population and Sample Size	Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)	Principal Findings
n=2068	and PAR to varying degrees by worksite.		predictability, absence from work, behavioural stress, somatic stress, emotional stress, and cognitive stress		workplaces. Method of analysis and results not detailed, but authors claim it is "statistically supported that": <ul style="list-style-type: none"> <li>the workplaces that did the most to improve the psychosocial work environment, achieved the highest drop in absence rate;</li> <li>the workplaces in which the psychosocial work environment, due to different reasons, became worse have experienced the highest increase in the absence rate;</li> <li>employees with reduced workability from workplaces where the psychosocial working environment was improved had a reduction in absence spells over the period;</li> <li>employees with reduced workability from workplaces where the psychosocial work environment was not improved, or was poor initially, had a considerable increase in absence spells over the period.</li> </ul>
Taris et al, 2003 <sup>99</sup> Domiciliary care employees from 81 organizations (The	Varying levels of Systems Approach across organisations	O, O/I, I to varying degrees by organisation.	All participating sites had intervention, but to varying degrees; unit of intervention and analysis = organisation; analysis by internal comparison relating intervention	3-***	Org-level: Overall means of job stress measures improved (job demands decreased, job control increased, social support increased, and emotional exhaustion decreased) over the

Intervention		Evaluation	
Systems Approach Rating (Low, Moderate, High) Additional Notes (PAR, NA/RA, OHS/HP) <sup>§</sup>	Intervention Level or Levels (E, O, I); Intervention Duration	Control or Comparison Groups; Evaluation Measures or outcomes	Study Design/Causal Inference Rating (3 stars to 5 stars) <sup>#</sup> Note if includes qualitative (QUAL)
<p><b>Study:</b> First Author, Year Population and Sample Size</p> <p>Netherlands) N = 26,563</p> <p>Includes NA/RA and PAR to varying degrees by organisation.</p>	<p>2 and 1/2 year follow-up evaluation</p>	<p>process to outcomes at individual and organisational levels.</p> <p>Measures of psychosocial working conditions: Emotional exhaustion (Maslach burnout inventory) as a measure of stress; job demands, job control, and social support;</p> <p>Type of intervention: work-directed (factual changes in work content and/or relations), person/work interface (to increase employee resistance to stressors), and person-directed (e.g., exercise, employee assistance programs, relaxation training)</p>	<p><b>Principal Findings</b></p> <p>intervention period (2 yrs).</p> <p>Relating intervention process to outcome at the organisational level showed:</p> <ul style="list-style-type: none"> <li>• Organisations usually implemented a wide variety of intervention activities;</li> <li>• The greater the number of interventions activities, the greater the improvement in job stress measures at individual level;</li> <li>• Work-directed, but not other, interventions were linked to job stress reduction at the individual level;</li> <li>• A higher number of work-directed interventions was significantly linked to larger decreases in job demands at the organisational level. Similar work-directed intervention improvement patterns were observed for other job stress measures at the org level, but these did not achieve statistical significance.</li> <li>• Overall conclusion is that work-directed, but not other,</li> </ul>

		Intervention		Evaluation	
<b>Study:</b> <b>First Author, Year</b> <b>Population and</b> <b>Sample Size</b>	<b>Systems</b> <b>Approach</b> <b>Rating (Low,</b> <b>Moderate,</b> <b>High)</b> <b>Additional</b> <b>Notes (PAR,</b> <b>NA/RA,</b> <b>OHS/HP)<sup>§</sup></b>	<b>Intervention</b> <b>Level or Levels</b> <b>(E, O, I);</b> <b>Intervention</b> <b>Duration</b>	<b>Control or Comparison Groups;</b> <b>Evaluation Measures or</b> <b>outcomes</b>	<b>Study</b> <b>Design/Causal</b> <b>Inference Rating</b> <b>(3 stars to 5</b> <b>stars)<sup>#</sup></b> <b>Note if includes</b> <b>qualitative</b> <b>(QUAL)</b>	<b>Principal Findings</b>
					interventions, are linked to job stress reduction, although the effect sizes were small.

## LITERATURE CITED

1. Adkins J, Quick J, Moe K. Building world-class performance in changing times. In: Murphy L, Cooper C, editors. *Healthy and Productive Work: An International Perspective*. London: Taylor & Francis; 2000. p. 107-131.
2. Aust B, Peter R, Siegrist J. Stress management in bus drivers: a pilot study based on the model of effort-reward imbalance. *International Journal of Stress Management* 1997;4(4):297-305.
3. Barrios CB, McCraty R, Cryer B. An inner quality approach to reducing stress and improving physical and emotional wellbeing at work. *Stress Medicine* 1997;13:193-201.
4. Bunce D, West MA. Stress management and innovation interventions at work. *Human Relations* 1996;49:209-232.
5. Cahill J. Computers and stress reduction in social service workers in New Jersey. *Conditions of Work Digest* 1992;11:197-203.
6. Cartwright S, Cooper C, Whatmore L. Improving communications and health in a government department. In: Murphy LR, Cooper CL, editors. *Healthy and Productive Work: An International Perspective*. London: Taylor & Francis.; 2000. p. 67-81.
7. Eriksson I, Moser V, Uden A, Orth-Gomer K. Using knowledge and discussion to decrease stress in Swedish public administration officials. *Conditions of Work Digest* 1992;11(2):214-219.
8. Griffin MA, Hart PM, Wilson-Evered E. Using employee opinion surveys to improve organisational health. In: Murphy LR, Cooper CL, editors. *Healthy and Productive Work: An International Perspective*. London: Taylor & Francis; 2000.
9. Griffiths A, Randall R, Santos A, Cox T. Senior nurses: Interventions to reduce work stress. In: Dollard M, Winefield A, Winefield H, editors. *Occupational Stress in the Service Professions*. London: Taylor & Francis; 2003. p. 169-191.
10. Innstrand ST, Espnes GA, Mykletun R. Job stress, burnout and job satisfaction: An intervention study for staff working with people with intellectual disabilities. *Journal of Applied Research in Intellectual Disabilities* 2004;17(2):119-126.
11. Israel BA, Schurman SJ, House JS. Action research on occupational stress: involving workers as researchers. *Int J Health Serv* 1989;19(1):135-155.
12. Israel BA, Schurman SJ, Hugentobler MK, House JS. A participatory action research approach to reducing occupational stress in the United States. *Conditions of Work Digest* 1992;11(2):152-163.
13. Heaney C, Israel BA, Schurman SJ, Baker E, House J, Hugentobler MK. Industrial relations, worksite stress reduction, and employee well-being: a participatory action research investigation. *Journal of Organisational Behavior* 1993;14:495-510.

14. Kalimo R, Toppinen S. Finland: organisational well-being. Ten years of research and development in a forest industry corporation. In: Kompier M, Cooper CL, editors. Preventing Stress, Improving Productivity: European Case Studies in the Workplace. London: Routledge; 1999.
15. Kawakami N, Araki S, Kawashima M, Masumoto T, Hayashi T. Effects of work-related stress reduction on depressive symptoms among Japanese blue-collar workers. *Scandinavian Journal of Work, Environment & Health* 1997;23(1):54-59.
16. Kvarnstrom S. From Taylorism to 1000 objective-orientated groups: experiences of a cultural revolution in an industrial concern. In: Cooper CL, Liukkonen P, Cartwright S, editors. Stress Prevention in the Workplace: Assessing the Costs and Benefits to Organisations. Dublin: European Foundation for the Improvement of Living and Working Conditions; 1996.
17. Lavoie-Tremblay M, Bourbonnais R, Viens C, Vezina M, Durand PJ, Rochette L. Improving the psychosocial work environment. *Journal of Advanced Nursing* 2005;49(6):655-664.
18. Logan MS, Ganster DC. An experimental evaluation of a control intervention to alleviate job-related stress. *Journal of Management* 2005;31(1):90-107.
19. Lourijsen E, Houtman L, Kompier MAJ, Grundemann R. The Netherlands: A hospital, "Healthy working for health". In: Kompier MAJ, Cooper CL, editors. Preventing stress, improving productivity: European case studies in the workplace. London: Routledge; 1999. p. 86-120.
20. Maes S, Verhoeven C, Kittel F, Scholten H. Effects of a Dutch wellness--health program: The Brabantia Project. *Am J Public Health* 1998;88(7):1037-1041.
21. Matrajt M. Using ergonomic analysis and group discussion to identify and prevent stress in managers and assembly-line workers. *Conditions of Work Digest* 1992;11(2):152-163.
22. Melchior ME, Philipsen H, Abu-Saad HH, Halfens RJ, van de Berg AA, Gassman P. The effectiveness of primary nursing on burnout among psychiatric nurses in long-stay settings. *Journal of Advanced Nursing* 1996;24(4):694-702.
23. Michie S, Wren B, Williams S. Reducing absenteeism in hospital cleaning staff: pilot of a theory based intervention. *Occupational and Environmental Medicine* 2004;61(4):345-349.
24. Mikkelsen A, Saksvik PØ, Landsbergis P. The impact of a participatory organizational intervention on job stress in community health care institutions. *Work & Stress* 2000;14(2):156-70.
25. Mikkelsen A, Saksvik PO. Impact of a participatory organizational intervention on job characteristics and job stress. *International Journal of Health Services* 1999;29(4):871-93.
26. Munz D, Kohler J, Greenberg C. Effectiveness of a comprehensive worksite management program: combining organisational and individual interventions. *International Journal of Stress Management* 2001;8(1):49-62.
27. Nijhuis FJ, Lendfers M, de Jong A, Janssen P, Ament A. Stress-related interventions in construction work. In: Cooper CL, Liukkonen P, Cartwright S, editors. Stress Prevention in the Workplace: Assessing the Costs and Benefits to



Organisations. Dublin: European Foundation for the Improvement of Living and Working Conditions; 1996.

28. Orth-Gomer K, Eriksson I, Moser V, Theorell T, Fredlund P. Lipid lowering through work stress reduction. *International Journal of Behavioral Medicine* 1994;1:204-214.
29. Poelmans S, Compernelle T, De Neve H, Buelens M, Rombouts J. Belgium: A pharmaceutical company. In: Kompier M, Cooper C, editors. *Preventing Stress, Improving Productivity: European Case Studies in the Workplace*. London: Routledge; 1999.
30. Sastry G. Using training to prevent or reduce stress in a coalmining company in India. *Conditions of Work Digest* 1992;11(2):268-275.
31. Schaubroeck J, Ganster DC, Sime W, Ditman D. A field experiment testing supervisory role clarification. *Personnel Psychology* 1993;46:1-25.
32. van Dierendonck D, Schaufeli WB, Buunk BP. The evaluation of an individual burnout intervention program: The role of inequity and social support. *Journal of Applied Psychology* 1998;83(3):392-407.
33. Kompier M, Cooper C. *Preventing Stress, Improving Productivity : European Case Studies in the Workplace*. London ; New York: Routledge; 1999.
34. Blomkvist V, Eriksen W, Theorell T, Ulrich R, Rasmanis G. Acoustics and psychosocial environment in intensive coronary care. *Occup Environ Med* 2005;62:e1.
35. Bond R, Bunce D. Job control mediates change in a work reorganization intervention for stress reduction. *Journal of Occupational Health Psychology* 2001;6:290-302.
36. de Croon EM, Kuijer PP, Broersen JP, Frings-Dresen MH. Information technology and road transport industry: how does IT affect the lorry driver? *Applied Ergonomics* 2004;35(4):313-20.
37. Elo AL, Leppanen A, Sillanpaa P. Applicability of survey feedback for an occupational health method in stress management. *Occupational Medicine (Oxford)* 1998;48(3):181-8.
38. Landsbergis PA, Vivona-Vaughan E. Evaluation of an occupational stress intervention in a public agency. *Journal of Organizational Behaviour* 1995;16:26-28.
39. Molleman E, Van Knippenberg A. Work redesign and the balance of control within a nursing context. *Human Relations* 1995;48(7):795-814.
40. Parkes KR. The effects of objective workload on cognitive performance in a field setting: a two-period cross-over trial. *Applied Cognitive Psychology* 1995;9:S153-S171.
41. Proctor R, Stratton-Powell H, Tarrier N. The impact of training and support on stress among care staff in nursing and residential homes for the elderly. *Journal of Mental Health* 1998;7(59-70).
42. Randall R, Griffiths A, Cox T. Evaluating organizational stress-management interventions using adapted study designs. *European Journal of Work and Organizational Psychology* 2005;14(1):23-41.

43. Reynolds S. Psychological well-being at work: is prevention better than cure? *Journal of Psychosomatic Research* 1997;43(1):93-102.
44. Rydstedt LW, Johansson G, Evans GW. The human side of the road: improving the working conditions of urban bus drivers. *Journal of Occupational Health Psychology*. 1998;3(2):161-71.
45. Evans GW, Johansson G, Rydstedt LW. Hassles on the job: a study of a job intervention with urban bus drivers. *Journal of Organizational Behaviour* 1999;20:199-208.
46. Smith M, Zehel D. A stress reduction intervention programme for meat processors emphasizing job design and work organization (United States). *Conditions of Work Digest* 1992;11(204-212).
47. Smoot SL, Gonzales JL. Cost-effective communication skills training for state hospital employees. *Psychiatric Services* 1995;46(8):819-22.
48. Terra N. The prevention of job stress by redesigning jobs and implementing self-regulating teams. In: Murphy L, Hurrell J, Sauter SL, Keita GP, editors. *Job Stress Interventions*. Washington: American Psychological Association; 1995.
49. Theorell T, Emdad R, Arnetz B, Weingarten AM. Employee effects of an educational program for managers at an insurance company. *Psychosomatic Medicine* 2001;63(5):724-33.
50. Wahlstedt K, Edling C. Organisational changes at a postal sorting terminal - their effects upon work satisfaction, psychosomatic complaints and sick leave. *Work and Stress* 1997;11(3):279-291.
51. Alexander C, GC S, Rainforth M, Carlisle T, Todd C, Oates J. Effects of the transcendental meditation program on stress reduction, health, and employee development: a prospective study in two occupational settings. *Anxiety, Stress and Coping* 1993;6:245-262.
52. Bond FW, Bunce D. Mediators of change in emotion-focused and problem-focused worksite stress management interventions. *Journal of Occupational Health Psychology* 2000;5(1):156-63.
53. Carson J, Cavagin J, Bunclark J, et al. Effective communication in mental health nurses: Did social support save the psychiatric nurse? *Nursing Times* 1999;4(1):31-42.
54. Cecil M, Foreman S. Effects of stress inoculation training and coworker support groups on teachers' stress. *Journal of School Psychology* 1990;28:105-118.
55. Cigrang JA, Todd SL, Carbone EG. Stress management training for military trainees returned to duty after a mental health evaluation: effect on graduation rates. *Journal of Occupational Health Psychology* 2000;5(1):48-55.
56. Cooper CL, Alliso T, Reynolds P, Sadri G. An individual-based counselling approach for combating stress in British Post Office employees. *Conditions of Work Digest* 1992;11(2):246-256.
57. Delvaux N, Razavi D, Marchal S, Bredart A, Farvacques C, Slachmuylder JL. Effects of a 105 hours psychological training program on attitudes, communication skills and occupational stress in oncology: a randomised study. *British Journal of Cancer* 2004;90(1):106-14.

58. Doctor R, Curtis D, Isaacs G. Psychiatric morbidity in policemen and the effect of brief psychotherapeutic intervention - a pilot study. *Stress Medicine* 1994;10:151-157.
59. Elliott T, Maples S. Stress management training for employees experiencing corporate acquisition. *Journal of Employment Counselling* 1991;28:107-114.
60. Eriksen HR, Ihlebaek C, Mikkelsen A, Gronningsaeter H, Sandal GM, Ursin H. Improving subjective health at the worksite: a randomized controlled trial of stress management training, physical exercise and an integrated health programme. *Occupational Medicine-Oxford* 2002;52(7):383-391.
61. Ewers P, Bradshaw T, McGovern J, Ewers B. Does training in psychosocial interventions reduce burnout rates in forensic nurses? *Journal of Advanced Nursing* 2002;37(5):470-6.
62. Francis ME, Pennebaker JW. Putting stress into words: the impact of writing on physiological, absentee, and self-reported emotional well-being measures. *American Journal of Health Promotion* 1992;6(4):280-7.
63. Freedy JR, Hobfoll SE. Stress inoculation for reduction of burnout: A conservation of resources approach. *Anxiety, Stress and Coping* 1994;6:311-325.
64. Gardiner M, Lovell G, Williamson P. Physician you can heal yourself! Cognitive behavioural training reduces stress in GPs. *Family Practice* 2004;21(5):545-551.
65. Goodspeed RB, DeLucia AG. Stress reduction at the worksite: an evaluation of two methods. *American Journal of Health Promotion* 1990;4(5):333-337.
66. Greco N. Individual-based training to reduce stress in managers and employees at a Canadian ministry. *Conditions of Work Digest* 1992;11(2):262-267.
67. Grossman AH, Silverstein C. Facilitating support groups for professionals working with people with AIDS. *Social Work* 1993;38(2):144-51.
68. Heron RJ, McKeown S, Tomenson JA, Teasdale EL. Study to evaluate the effectiveness of stress management workshops on response to general and occupational measures of stress. *Occupational Medicine (Oxford)* 1999;49(7):451-7.
69. Hyman RB. Evaluation of an intervention for staff in a long-term care facility using a retrospective pretest design. *Evaluation & the Health Professions* 1993;16(2):212-24.
70. Iwi D, Watson J, Barber P, Kimber N, Sharman G. The self-reported well-being of employees facing organizational change: effects of an intervention. *Occupational Medicine (Oxford)* 1998;48(6):361-8.
71. Jenkins S, Calhoun J. Teacher stress: issues and intervention. *Psychology in the Schools* 1991;28(60-70):???
72. Johanning E, Landsbergis P, Geissler H, Karazmann R. Cardiovascular risk and back disorder intervention study of mass transit operators. *International Journal of Occupational and Environmental Health* 1996;2:79-87.
73. Kushnir T, Malkinson R. A rational-emotive group intervention for preventing and coping with stress among safety officers. *Journal of Rational-Emotive & Cognitive-Behavior Therapy* 1993;11(4):195-206.

74. Le Scanff C, Taugis J. Stress management for police special forces. *Journal of Applied Sport Psychology* 2002;14(4):330-343.
75. Lee S, Crockett MS. Effect of assertiveness training on levels of stress and assertiveness experienced by nurses in Taiwan, Republic of China. *Issues in Mental Health Nursing*. 1994;15:419-432.
76. McCue JD, Sachs CL. A stress management workshop improves residents' coping skills. *Archives of Internal Medicine* 1991;151(11):2273-7.
77. Meier A. Offering social support via the Internet: A case study of an online support group for social workers. *Journal of Technology in Human Services* 2000;17(2/3):237-66.
78. Michie S. Reducing absenteeism by stress management: Valuation of stress counselling service. *Work and Stress* 1996;10(4):367-372.
79. Pelletier KR, Rodenburg A, Vinther A, Chikamoto Y, King AC, Farquhar JW. Managing job strain: a randomized, controlled trial of an intervention conducted by mail and telephone. *Journal of Occupational & Environmental Medicine* 1999;41(4):216-23.
80. Peters K, Carlson J. Worksite stress management with high-risk maintenance workers: a controlled study. *International Journal of Stress Management* 1999;6(1):21-45.
81. Pruitt RH. Effectiveness and cost efficiency of interventions in health promotion. *Journal of Advanced Nursing* 1992;17(8):926-32.
82. Reynolds S, Taylor E, Shapiro D. Session impact and outcome in stress management training. *Journal of Community and Applied Social Psychology* 1993;3:325-337.
83. Robinson RC, Mitchell JT. Evaluation of psychological debriefings. *Journal of Traumatic Stress* 1993;6:367-382.
84. Schaufeli WB. The evaluation of a burnout workshop for community nurses. *Journal of Health & Human Services Administration* 1995;18(1):11-30.
85. Sheppard W, Staggers Jr F, John L. The effects of a stress management program in a high security governmental agency. *Anxiety, Stress and Coping* 1997;10:341-350.
86. Taylor MW. Effects of initial stress level, social support and participation in an exercise or music condition on the posttreatment stress, depression, and anxiety of nurses.: ED. D. St John's University; 1991.
87. Teasdale EL, Heron RJL, Tomenson JA. Bringing health to life. In: Murphy LR, Cooper CL, editors. *Healthy and productive work: An international perspective*. London: Taylor & Francis; 2000.
88. Toivanen H, Helin P, Hanninen O. Impact of regular relaxation training and psychosocial working factors on neck-shoulder tension and absenteeism in hospital cleaners. *Journal of Occupational Medicine* 1993;35(11):1123-30.
89. Toivanen H, Lansimies E, Jokela V, Hanninen O. Impact of regular relaxation training on the cardiac autonomic nervous system of hospital cleaners and bank employees. *Scandinavian Journal of Work, Environment & Health* 1993;19(5):319-325.

90. Tsai S, Crockett M. Effects of relaxation training, combining imagery, and meditation on the stress level of Chinese nurses working in modern hospitals in Taiwan. *Issues in Mental Health Nursing* 1993;14(1):51-66.
91. Vines SW. Relaxation with guided imagery: effects on employees' psychological distress and health seeking behaviors. *American Association of Occupational Health Nursing Journal* 1994;42(5):206-13.
92. Whatmore L, Cartwright S, Cooper C. United Kingdom: evaluation of a stress management programme in the public sector. In: Kompier M, Cooper CL, editors. *Preventing Stress, Improving Productivity: European Case Studies in the Workplace*. London: Routledge; 1999. p. 149-174.
93. Wilholm C, Arnetz B, Berg M. The impact of stress management on computer-related skin problems. *Stress Medicine* 2000;16(279-285).
94. Eklof M, Ingelgard A, Hagberg M. Is participative ergonomics associated with better working environment and health? A study among Swedish white-collar VDU users. *International Journal of Industrial Ergonomics* 2004;34(5):355-366.
95. Eklof M, Hagberg M, Toomingas A, Tornqvist EW. Feedback of workplace data to individual workers, workgroups or supervisors as a way to stimulate working environment activity: a cluster randomized controlled study. *International Archives of Occupational and Environmental Health* 2004;77(7):505-514.
96. Lindstrom K, Schrey K, Ahonen G, Kaleva S. The effects of promoting organisational health on worker well-being and organisational effectiveness in small and medium-sized enterprises. In: Murphy L, Cooper CL, editors. *Healthy and Productive Work: An International Perspective*. London: Taylor & Francis; 2000.
97. Nielsen M, Kristensen T, Smith-Hansen L. The Intervention Project on Absence and Well-being (IPAW): Design and results from the baseline of a 5-year study. *Work and Stress* 2002;16(3):191-206.
98. Nielsen M. The intervention project on absence and well-being (IPAW)--Denmark. In: EASHW, editor. *How to Tackle Psychosocial Issues and Reduce Work-Related Stress*. Luxembourg: European Agency for Safety and Health at Work (EASHW); 2002. p. 52-58.
99. Taris T, Kompier M, Geurts A, et al. Stress management interventions in the Dutch domiciliary care sector: findings from 81 organizations. *International Journal of Stress Management* 2003;10(4):297-325.